arXiv:1702.06281v1 [math.CO] 21 Feb 2017

THE NUMBERS OF EDGES OF 5-POLYTOPES

TAKUYA KUSUNOKI AND SATOSHI MURAI

ABSTRACT. A basic combinatorial invariant of a convex polytope P is its f-vector $f(P) = (f_0, f_1, \ldots, f_{\dim P-1})$, where f_i is the number of i-dimensional faces of P. Steinitz characterized all possible f-vectors of 3-polytopes and Grünbaum characterized the first two entries of the f-vectors of 4-polytopes. In this paper, we characterize the first two entries of the f-vectors of 5-polytopes.

1. INTRODUCTION

The study of f-vectors of convex polytopes is one of the central research topic in convex geometry. We call a d-dimensional convex polytope a *d-polytope*. For a convex polytope (or a polyhedral complex) P, we write $f_i(P)$ for the number of i-dimensional faces of P. The *f-vector* of a d-polytope P is the vector $f(P) = (f_0(P), f_1(P), \ldots, f_{d-1}(P))$. In 1906, Steinitz characterized all possible f-vectors of 3-polytopes (see [Gr, §10.3]). While a characterization of f-vectors of 4-polytopes is a big open problem in convex geometry, for any $0 \leq i < j \leq 3$, the following set was characterized by Grünbaum [Gr], Barnette [Ba] and Barnette–Reay [BR] (see also [BL, Theorem 3.9])

$$\big\{(f_i(P), f_j(P)) : P \text{ is a 4-polytope}\big\}.$$

Moreover, Sjöberg and Ziegler [SZ] recently characterize all possible values of the pairs (f_0, f_{03}) of flag face numbers of 4-polytopes. In this paper, we characterize all possible (f_0, f_1) pairs of 5-polytopes.

Let

$$\mathcal{E}^d = \{(f_0(P), f_1(P)) : \ P \text{ is a } d\text{-polytope}\}.$$

The set $\mathcal{E}^3$ was determined by Steinitz in 1906 who shows that

$$\mathcal{E}^3 = \{(v, e) : \tfrac{3}{2}v \leq e \leq 3v - 6\}.$$

Note that, by Euler's relation, this actually determines all possible f-vectors of 3-polytopes. In higher dimensions, it is easy to see that any d-polytope P satisfies

$$\frac{d}{2} f_0(P) \leq f_1(P) \leq \binom{f_0(P)}{2}. \tag{1}$$

Indeed, the first inequality follows since $f_1(P)$ equals to $\frac{1}{2}$ times the sum of degrees of the vertices of P and since each vertex of P has degree $\geq d$. Grünbaum [Gr, §10.4] proved that the inequality (1) characterizes $\mathcal{E}^4$, with four exceptions. More precisely, he proved the following statement.

The second author was partially supported by KAKENHI16K05102.

Theorem 1.1 (Grünbaum)**.**

$$\mathcal{E}^4 = \left\{(v,e) : 2v \le e \le \binom{v}{2}\right\} \setminus \big\{(6,12),(7,14),(8,17),(10,20)\big\}.$$

In dimension 5, the situation is more complicated. The set $\mathcal{E}^5$ is close to the set of integer points satisfying (1), but there are not only a finite list of exceptions but also an infinite family of exceptions. Indeed, our main result is the following.

Theorem 1.2. *Let* $L = \{(v, \lfloor \frac{5}{2}v + 1 \rfloor) : v \ge 6\}$ *and* $G = \{(8,20),(9,25),(13,35)\}$. *Then*

$$\mathcal{E}^5 = \left\{(v,e) : \frac{5}{2}v \le e \le \binom{v}{2}\right\} \setminus (L \cup G).$$

Here $\lfloor a \rfloor$ denotes the integer part of a rational number a. Note that it is not hard to see $(8,20) \notin \mathcal{E}^5$ since a 5-polytope P with $f_1(P) = \frac{5}{2}f_0(P)$ must be a simple polytope. Also, $(9,25) \notin \mathcal{E}^5$ was proved in [PUY] recently. The following Table 1 illustrates the shape of $\mathcal{E}^5$. In the table, we place the possible numbers of edges for a fixed number of vertices in each column so that the bottom number is $\lfloor \frac{5}{2}v + \frac{1}{2} \rfloor$, where the symbols '$L$' and '$G$' represent the positions of elements of L and G respectively. (For example, the table says that if the number of the vertices is 9, then the possible numbers of edges are $24, 26, 27, \ldots, 36$.)

e													
					45	⋮	⋮	⋮	⋮	⋮	⋮	⋮	⋮
				36	⋮	⋮	⋮	⋮	⋮	⋮	⋮	⋮	⋮
			28	⋮	⋮	⋮	⋮	⋮	⋮	⋮	⋮	⋮	⋮
			⋮	⋮	⋮	⋮	⋮	⋮	⋮	⋮	⋮	⋮	⋮
		21	23	26	28	31	33	36	38	41	43	46	48
		20	22	G	27	30	32	G	37	40	42	45	47
		19	L	24	L	29	L	34	L	39	L	44	L
	15	L	G	L	25	L	30	L	35	L	40	L	45
	6	7	8	9	10	11	12	13	14	15	16	17	18

v

Table 1: The table of $\mathcal{E}^5$

It would be interesting to determine $\mathcal{E}^d$ for $d \ge 6$, and more generally to characterize the set $\{(f_i(P), f_j(P)) : P \text{ is a } d\text{-polytope}\}$ for any $0 \le i < j < d$. About the latter problem, Sjöberg and Ziegler [SZ] recently study the case when $i = 0$ and $j = d - 1$.

2. SUFFICIENCY

In this section, we prove the sufficiency part of Theorem 1.2. If a polytope Q is the pyramid over a polytope P, then we have

$$f_0(Q) = f_0(P) + 1 \text{ and } f_1(Q) = f_0(P) + f_1(P).$$

This simple fact and Theorem 1.1 prove the next lemma.

Lemma 2.1.

$$\mathcal{E}^5 \supset \left\{(v,e) : 3v-3 \le e \le \binom{v}{2}\right\} \setminus \big\{(7,18),(8,21),(9,25),(11,30)\big\}.$$

Let P be a d-polytope. The *degree* $\deg v$ of a vertex v of P is the number of edges of P that contain v. We say that a vertex v is *simple* if $\deg v = d$. Let $V(P)$ be the vertex set of P.

Lemma 2.2. *If P is a 5-polytope such that $f_1(P) \le 3f_0(P)-1$, then P has a simple vertex.*

Proof. Observe $\deg v \ge 5$ for any $v \in V(P)$. Since $\frac{1}{2}\sum_{v\in V(P)} \deg v = f_1(P) = 3f_0(P)-1$, there must exist a vertex $v \in V(P)$ of degree < 6. □

Let

$$X = \big\{(v,e) : \tfrac{5}{2}v \le e \le \tbinom{v}{2}\big\} \setminus (L \cup G)$$

be the right-hand side of Theorem 1.2, and let

$$X_k = \{(v,e) \in X : v = k\}$$

for $k \in \mathbb{Z}$. We want to prove $\mathcal{E}^5 \supset X_k$ for all $k \ge 6$. To prove this, we use truncations. For a 5-polytope P and its vertex $v \in V(P)$, we write $\mathrm{tr}(P,v)$ for a polytope obtained from P by truncating the vertex v. If v is simple, then

$$f_0(\mathrm{tr}(P,v)) = f_0(P)+4 \text{ and } f_1(\mathrm{tr}(P,v)) = f_1(P)+10.$$

Lemma 2.3. *For $k \ge 6$ with $k \notin \{8,9,13\}$, if $\mathcal{E}^5 \supset X_k$ then $\mathcal{E}^5 \supset X_{k+4}$.*

Proof. Since $k \notin \{8,9,13\}$,

$$X_k = \big\{(k,e) : \tfrac{5}{2}k \le e \le \tbinom{k}{2}\big\} \setminus \big\{\big(k, \lfloor \tfrac{5}{2}k+1 \rfloor\big)\big\}.$$

By Lemma 2.2, for any 5-polytope P with $f_0(P) = k$ and $f_1(P) \le 3k-1$, we can make a 5-polytope Q with

$$f_0(Q) = k+4 \text{ and } f_1(Q) = f_1(P)+10$$

by truncating a simple vertex from P. Since $\mathcal{E}^5 \supset X_k$, this implies

$$\begin{aligned}\mathcal{E}^5 &\supset \left\{(k+4,e+10) : \frac{5}{2}k \le e \le 3k-1\right\} \setminus \left\{\left(k+4, \left\lfloor \frac{5}{2}k+11 \right\rfloor\right)\right\} \\ &= \left\{\big((k+4),e'\big) : \frac{5}{2}(k+4) \le e' \le 3(k+4)-3\right\} \setminus \left\{\left(k+4, \left\lfloor \frac{5}{2}(k+4)+1 \right\rfloor\right)\right\}.\end{aligned}$$

The above inclusion and Lemma 2.1 prove the desired statement. □

Now, we prove the main result of this section. For a convex polytope P, we write P^* for its dual polytope. In the rest of the paper, if a face of a convex polytope is a simplex, then we call it a *simplex face*. A face which is not a simplex is called a *non-simplex face*.

Theorem 2.4. $\mathcal{E}^5 \supset X$.

Proof. By Lemma 2.3, it is enough to show that

$$\mathcal{E}^5 \supset X_6 \cup X_7 \cup X_8 \cup X_9 \cup X_{12} \cup X_{13} \cup X_{17}. \tag{2}$$

Let $\varphi(v) = 3v - 3$. By Lemma 2.1, $(v, e) \in \mathcal{E}^5$ if $\varphi(v) \leq e \leq \binom{v}{2}$ and $(v, e) \notin \{(7, 18), (8, 21), (9, 25), (11, 30)\}$. Observe $\varphi(6) = 15, \varphi(7) = 18, \varphi(8) = 21, \varphi(9) = 24, \varphi(12) = 33, \varphi(13) = 36, \varphi(17) = 48$. Then, to prove (2), what we must prove is

$$\mathcal{E}^5 \supset \{(12, 30), (12, 32), (13, 34), (17, 44), (17, 45), (17, 46), (17, 47)\}. \tag{3}$$

(See also Table 1.) Note that this observation says $\mathcal{E}^5 \supset X_k$ for $k \leq 9$.

Let C be the cyclic 5-polytope with 7 vertices. Then $f(C) = (7, 21, 34, 30, 12)$ (see [Br, §18]). Hence $f_0(C^*) = 12$ and $f_1(C^*) = 30$, and therefore $(12, 30) \in \mathcal{E}^5$. Let C' be the polytope obtained from C^* by truncating its vertex. Note that every vertex of C^* is simple. Since a truncation of a simple vertex creates a simplex facet, C' contains a simplex facet F. Let C'' be the polytope obtained from C' by adding a pyramid over F. Then

$$f_0(C'') = f_0(C') + 1 = f_0(C^*) + 5 = 17$$

and

$$f_1(C'') = f_1(C') + 5 = f_1(C^*) + 15 = 45.$$

Hence $(17, 45) \in \mathcal{E}^5$. We already see $(8, 22), (9, 24) \in \mathcal{E}^5$. Then, using truncations of simple vertices and Lemma 2.2, we have $(12, 32), (13, 34), (17, 44) \in \mathcal{E}^5$. Also, since $(13, 36), (13, 37) \in \mathcal{E}^5$, by the same argument we have $(17, 46), (17, 47) \in \mathcal{E}^5$. These complete the proof of the theorem. □

3. Necessity

In this section, we prove the necessity part of Theorem 1.2. We first show that any element in L is not contained in $\mathcal{E}^5$. We introduce some lemmas which we need. The following fact appears in [Gr, §6.1] (see also [Zi, Problem 6.8]).

Lemma 3.1. *There are exactly four combinatorially different* 4*-polytopes with* 6 *facets. They are*

- (P_A) *Pyramid over a square pyramid;*
- (P_B) *Pyramid over a triangular prism;*
- (P_C) *A polytope obtained from a* 4*-simplex by truncating its vertex;*
- (P_D) *Product of two triangles.*

Here are Schlegel diagrams and a list of facets of P_A, P_B, P_C and P_D.



P_A P_B P_C P_D

Type	Facets
P_A	two square pyramids, four tetrahedra
P_B	three square pyramids, one triangular prism, two tetrahedra
P_C	four triangular prisms, two tetrahedra
P_D	six triangular prisms

Recall that a convex polytope P is said to be *simplicial* if all its proper faces are simplices. A *simplicial k-sphere* is a simplicial complex whose geometric realization is homeomorphic to the k-sphere. The boundary complex of a simplicial d-polytope is a simplicial $(d-1)$-sphere. The next statement easily follows from the Lower Bound Theorem (see [Ka]) and the Upper Bound Theorem (see [St, Corollary II.3.5]) for simplicial spheres.

Lemma 3.2. *Let Δ be a simplicial 3-sphere.*

(i) $f_3(\Delta) \neq 6, 7, 10$.
(ii) *If $f_3(\Delta) = 9$, then Δ is neighbourly, that is, every pair of vertices of Δ are connected by an edge.*

Proof. By the Lower Bound Theorem and the Upper Bound Theorem, we have

- if $f_0(\Delta) = 5$, then $5 \leq f_3(\Delta) \leq 5$;
- if $f_0(\Delta) = 6$, then $8 \leq f_3(\Delta) \leq 9$;
- if $f_0(\Delta) \geq 7$, then $11 \leq f_3(\Delta)$.

These clearly imply (i). The statement (ii) follows from the fact that if the number of facets of a simplicial $(d-1)$-sphere equals to the bound in the Upper Bound Theorem, then it must be neighbourly (see e.g. the proof of [Br, Theorem 18.1]). □

We now prove that any element in L is not contained in $\mathcal{E}^5$.

Proposition 3.3. *If P is a 5-polytope, then $f_1(P) \neq \lfloor \frac{5}{2} f_0(P) + 1 \rfloor$.*

Proof. Suppose to the contrary that $f_1(P) = \lfloor \frac{5}{2} f_0(P) + 1 \rfloor$. We first consider the case when $f_0(P)$ is odd. Then, since $\sum_{v \in V(P)} \deg v = 2f_1(P) = 5f_0(P) + 1$, P has one vertex having degree 6 and all other vertices have degree 5. Then P^* has one facet F with $f_3(F) = 6$ and all other facets of P^* are simplices. However this implies that the 4-polytope F must be simplicial, which contradicts Lemma 3.2(i).

Next, we consider the case when $f_0(P)$ is even. In this case, $\sum_{v \in V(P)} \deg v = 2f_1(P) = 5f_0(P) + 2$, so one of the following two cases occurs:

(a) P^* has one facet F with $f_3(P) = 7$ and all other facets of P^* are simplices;
(b) P^* has two facets F and G with $f_3(F) = f_3(G) = 6$ and all other facets of P^* are simplices.

Since there are no simplicial 4-polytope with 7 facets by Lemma 2.2(i), the case (a) cannot occur. Also, if the case (b) occurs, then F and G can have at most one non-simplex facet. However, Lemma 3.1 says that any 4-polytope with 6 facets have at least two non-simplex facets. □

Next, we show that any element of G is not contained in $\mathcal{E}^5$. Let $\phi(v,d) = \frac{1}{2}dv + \frac{1}{2}(v-d-1)(2d-v)$. The following result was proved by Pineda-Villavicencio, Udon and Yost [PUY, Theorems 6 and 19].

Theorem 3.4. *Let P be a d-polytope.*

(i) *If $f_0(P) \leq 2d$, then $f_1(P) \geq \phi(f_0(P), d)$.*
(ii) *If $d \geq 4$, then $(f_0(P), f_1(P)) \neq (d+4, \phi(d+4,d)+1)$.*

By considering the special case when $d = 5$ of the above theorem, we obtain the following.

Corollary 3.5. $(8,20), (9,25) \notin \mathcal{E}^5$.

By Proposition 3.3 and Corollary 3.5, to prove Theorem 1.2, we only need to prove $(13,35) \notin \mathcal{E}^5$. We will prove this in the rest of this paper.

Let P be a 5-polytope. For faces $F_1, F_2, \ldots, F_k$ of P, we write $\langle F_1, \ldots, F_k \rangle$ for the polyhedral complex generated by $F_1, \ldots, F_k$. Let $\{G_1, \ldots, G_l\}$ be a subset of the set of facets of P. Then any 3-face of $\Gamma = \langle G_1, \ldots, G_l \rangle$ is contained in at most two facets of Γ. We write

$$\partial\Gamma = \langle H \in \Gamma : \ H \text{ is a 3-face of } \Gamma \text{ contained in exactly one facet of } \Gamma \rangle.$$

We often use the following trivial observation: If $\{G_1, \ldots, G_l\}$ is the set of non-simplex facets of P, then $\partial\langle G_1, \ldots, G_l \rangle$ is a simplicial complex.

We say that a d-polytope P is *almost simplicial* if all facets of P except for one facet are simplices. (We consider that simplicial polytopes are not almost simplicial.) The next lemma can be checked by using a complete list of 4-polytopes with at most 8 vertices (see [FMM]), but we write its proof for completeness.

Lemma 3.6. *Let P be a 4-polytope.*

(i) *Suppose that P is almost simplicial and $f_3(P) = 7$. Then P is the pyramid over a triangular bipyramid.*
(ii) *Suppose that P is almost simplicial and $f_3(P) = 8$. Then P does not contain a triangular bipyramid as a facet.*
(iii) *Suppose that $f_3(P) = 7$ and P has exactly two non-simplex facets F and G. Then none of F and G are square pyramids.*

Proof. (i) Let F be the unique non-simplex facet of P. Clearly, F is simplicial and $f_2(F) \leq f_3(P) - 1 = 6$ since, for each 2-face of F, there is a unique 3-face of P that contains it other than F. Since a 3-simplex and a triangular bipyramid are the only simplicial 3-polytopes having at most 6 facets, F is a triangular bipyramid. Then, since P has 7 facets, P must be the pyramid over F.

(ii) Let F be the unique non-simplex facet of P. If F is a triangular bipyramid, then by subdividing F into two tetrahedra without introducing edges, one obtains a simplicial 3-sphere Δ with 9 facets. Since F is a triangular bipyramid, there are two vertices u and v of F such that u and v are not connected by an edge in F. These vertices are not connected by an edge in Δ by the construction of Δ, which contradicts Lemma 3.2(ii) saying that Δ must be neighbourly.

(iii) Suppose to the contrary that F is a square pyramid. We claim that G is also a square pyramid. Indeed, since $\partial\langle F, G\rangle$ is a simplicial complex, G contains exactly one non-simplex facet, and this facet must be a square and equals to $F \cap G$. Also,

$$f_2(G) \leq f_3(P) - 1 = 6.$$

Let Δ be a simplicial 2-sphere obtained from G by subdividing the square $F \cap G$ into two triangles. Then $f_2(\Delta) \leq 7$, but since the number of 2-faces of a simplicial 2-sphere is even, $f_2(\Delta) = 6$ and therefore $f_2(G) = 5$. This forces that G is a square pyramid.

Let $F = \operatorname{conv}(v_1, a, b, c, d)$ and $G = \operatorname{conv}(v_2, a, b, c, d)$, where $\operatorname{conv}(v_1, \ldots, v_k)$ denotes the convex hull of points $v_1, \ldots, v_k$. Note that $\operatorname{conv}(a, b, c, d)$ is a square. We assume that $\operatorname{conv}(a, c)$ and $\operatorname{conv}(b, d)$ are non-edges of P. By subdividing each F and G into two tetrahedra by adding an edge $\operatorname{conv}(a, c)$, we can make a simplicial 3-sphere Γ with $f_3(\Gamma) = f_3(P) + 2 = 9$. By the construction of Γ, $\operatorname{conv}(b, d)$ is not an edge of Γ, but this contradicts Lemma 3.2(ii). □

We also recall some known results on h-vectors of simplicial balls and their boundaries. For a simplicial complex Δ of dimension $d-1$, its *h-vector* $h(\Delta) = (h_0(\Delta), h_1(\Delta), \ldots, h_d(\Delta))$ is defined by

$$h_i(\Delta) = \sum_{j=0}^{i} (-1)^{i-j} \binom{d-j}{i-j} f_{j-1}(\Delta)$$

where $f_{-1}(\Delta) = 1$. A *simplicial d-ball* is a simplicial complex whose geometric career is homeomorphic to a d-dimensional ball. The following facts are known. See [St, Chapter II and Problem 12].

Lemma 3.7. *Let Δ be a simplicial d-ball and $(h_0, h_1, \ldots, h_{d+1})$ its h-vector. Then*

(i) $h_0 = 1$ *and* $h_{d+1} = 0$.
(ii) $h_0 + \cdots + h_{d+1} = f_d(\Delta)$.
(iii) $h_i \geq 0$ *for all* i.
(iv) *The h-vector of the boundary complex of Δ is*

$$(h_0, h_0 + h_1 - h_d, h_0 + h_1 + h_2 - h_d - h_{d-1}, \cdots).$$

We now complete the proof of Theorem 1.2.

Theorem 3.8. $(13, 35) \notin \mathcal{E}^5$.

Proof. Suppose to the contrary that there is a 5-polytope P such that $f_0(P) = 13$ and $f_1(P) = 35$. Let $v_1, \ldots, v_{13}$ be the vertices of P with $\deg v_1 \geq \cdots \geq \deg v_{13}$ and let $D = (\deg v_1, \deg v_2, \ldots, \deg v_{13})$. Since

$$\sum_{k=1}^{13} \deg v_k = 2f_1(P) = 70$$

and $\deg v_k \geq 5$ for each k, D must be one of the following.

(1) $D = (10, 5, \ldots, 5)$;
(2) $D = (9, 6, 5, \ldots, 5)$;

(3) $D = (8, 7, 5, \ldots, 5)$;
(4) $D = (8, 6, 6, 5, \ldots, 5)$;
(5) $D = (7, 7, 6, 5, \ldots, 5)$;
(6) $D = (7, 6, 6, 6, 5, \ldots, 5)$;
(7) $D = (6, 6, 6, 6, 6, 5, \ldots, 5)$.

Below we will show a contradiction for each case.

(1) Suppose $D = (10, 5, \ldots, 5)$. Then P^* has a 4-face F with $f_3(F) = 10$. This F must be a simplicial 4-polytope since all other facets of P^* are simplices, which contradicts Lemma 3.2(i) saying that there are no simplicial 4-polytopes with 10 facets.

(2) Suppose $D = (9, 6, 5, \ldots, 5)$. P^* has only two non-simplex 4-faces F and G. These 4-faces can have at most one non-simplex 3-face. By the assumption on D, F or G must have 6 facets. This contradicts Lemma 3.1 saying that any 4-polytope with 6 facets has at least two non-simplex facets.

(3) Suppose $D = (8, 7, 5, \ldots, 5)$. Let F and G be the 4-faces of P^* with $f_3(F) = 8$ and $f_3(G) = 7$. Then G is not simplicial by Lemma 3.2(i) and therefore F is also not simplicial. Hence F and G are almost simplicial and $F \cap G$ is a 3-polytope which is not a simplex. Since $f_3(F) = 8$ and $f_3(G) = 7$, this contradicts Lemma 3.6(i) and (ii).

(4) Suppose $D = (8, 6, 6, 5, \ldots, 5)$. Let F, G and G' be 4-faces of P^* with $f_3(F) = 8$ and $f_3(G) = f_3(G') = 6$. Since $\partial\langle F, G, G'\rangle$ is a simplicial complex, the polytopes F, G and G' have at most two non-simplex 3-faces. By Lemma 3.1, G and G' must be the polytope P_A, and F must have 2 square pyramids as its 3-faces. Thus, the 4-polytope F has 6 tetrahedra and 2 square pyramids as its facets. This implies

$$f_0(F^*) = f_3(F) = 8 \quad \text{and} \quad f_1(F^*) = f_2(F) = \frac{1}{2}(5 \times 2 + 4 \times 6) = 17.$$

However, this contradicts Theorem 1.1 saying that $(8, 17) \notin \mathcal{E}^4$.

(5) Suppose $D = (7, 7, 6, 5, \ldots, 5)$. Let F, F' and G be 4-faces of P^* with $f_3(F) = f_3(F') = 7$ and $F(G) = 6$. Since all other 4-faces of P^* are simplices, F, F' and G have at most two non-simplex 3-faces. Then, by Lemma 3.1, G must be the polytope P_A, and F and F' have a square pyramid as its facets. By Lemma 3.6(i), F and F' are not almost simplicial. Hence F and F' have exactly two non-simplex facets, but this contradicts Lemma 3.6(iii).

(6) Suppose $D = (7, 6, 6, 6, 5, \ldots, 5)$. Let F, G, G' and G'' be 4-facets of P^* with $f_3(F) = 7$ and $f_3(G) = f_3(G') = f_3(G'') = 6$. Since all other 4-faces of P^* are simplices, each of F, G, G' and G'' can have at most three non-simplex 3-faces. By Lemma 3.1, G, G' and G'' must be the polytope P_A and have exactly two non-simplex 3-faces. Since $\partial\langle F, G, G', G''\rangle$ is a simplicial complex, one of the following situations must occur:

(a) F is a simplicial polytope;
(b) F has exactly two square pyramids as its facets and all other facets are simplices.

However, (a) cannot occur by Lemma 3.2(i) and (b) cannot occur by Lemma 3.6(iii).

(7) Suppose $D = (6, 6, 6, 6, 6, 5, \ldots, 5)$. Let $F_1, \ldots, F_5$ be the 4-faces of P^* with $f_3(F_1) = \cdots = f_3(F_5) = 6$. Observing that each of $F_1, \ldots, F_5$ must be one of P_A, P_B, P_C and P_D in Lemma 3.1. It is not hard to see that one of the following situations must occur;

(a) All the $F_1, \ldots, F_5$ are P_A;
(b) All the $F_1, \ldots, F_5$ are P_C;
(c) F_1 and F_2 are P_B. F_3, F_4 and F_5 are P_A.

We first show that (a) cannot occur. We may assume that $F_i \cap F_{i+1}$ is a square pyramid for $i = 1, 2, \ldots, 5$, where $F_6 = F_0$. Since P_A has only one square as its 2-faces, $K = F_1 \cap F_2 \cap \cdots \cap F_5$ must be this square, and the polyhedral complex generated by the facets of P^* that do not contain K is a shellable simplicial 4-ball by a line shelling [Zi, §8.2]. Let B be this ball. Clearly,

$$B = \langle G : G \text{ is a simplex facet of } P^* \rangle.$$

Hence B has $13 - 5 = 8$ facets. Also, since each $F_i \cap F_{i+1}$ is a pyramid over the square K, the faces $F_1, \ldots, F_5$ can be written as

$$F_1 = \mathrm{conv}(v_1, v_2, u_1, u_2, u_3, u_4),$$

$$F_2 = \mathrm{conv}(v_2, v_3, u_1, u_2, u_3, u_4),$$

$$\vdots$$

$$F_5 = \mathrm{conv}(v_5, v_1, u_1, u_2, u_3, u_4)$$

with $K = \mathrm{conv}(u_1, \ldots, u_4)$. Then it follows that $\Gamma = \partial\langle F_1, \ldots, F_5 \rangle$ is the join of the 5-cycle and the 4-cycle, and its h-vector is

$$(1, 5, 8, 5, 1)$$

since its entries coincide with the coefficients of the polynomial $(1+3t+t^2)(1+2t+t^2)$. Let $h(B) = (1, h_1, h_2, h_3, h_4, 0)$. Lemmas 3.7 and 3.8 say

$$1 + h_1 - h_4 = 5,\ 1 + h_1 + h_2 - h_3 - h_4 = 8 \text{ and } 1 + h_1 + h_2 + h_3 + h_4 = 8.$$

Then it is easy to see $h(B) = (1, 4, 3, 0, 0, 0)$. Let $G_1, \ldots, G_8$ be a shelling of B. Let

$$R_j = \{v \in V(G_j) : \mathrm{conv}(V(G_j) \setminus \{v\}) \in \langle G_1, \ldots, G_{j-1} \rangle\}$$

and $S_j = V(G_j) \setminus R_j$, where $V(G_j)$ is the vertex set of G_j. Then

$$h_i = h_i(B) = |\{j : |R_j| = i\}|$$

for all i (see [Zi, §8.3]). Since $h_3 = h_4 = 0$, we have $|S_j| = 5 - |R_j| \geq 3$ for all j. By the definition of a shelling, $\mathrm{conv}(S_8)$ is a missing face of ∂B, that is, $\mathrm{conv}(S_8)$ is not a face of ∂B but any its proper face is a face of ∂B. Thus ∂B has a missing face of dimension ≥ 2. However, the join of two cycles of length ≥ 4 does not have any missing face of dimension ≥ 2, and $\partial B = \partial\langle F_1, \ldots, F_5 \rangle$ is the join of the 5-cycle and the 4-cycle, a contradiction.

We next prove that (b) cannot occur. Observe that P_C has 4 non-simplex facets. Since $\partial\langle F_1, \ldots, F_5\rangle$ is a simplicial complex, each $F_i \cap F_j$ must be a triangular prism. Then it is easy to see that we can write

$$\begin{aligned}
F_1 &= \mathrm{conv}(x_1, x_2, x_3, x_4, y_1, y_2, y_3, y_4),\\
F_2 &= \mathrm{conv}(x_1, x_2, x_3, x_5, y_1, y_2, y_3, y_5),\\
F_3 &= \mathrm{conv}(x_1, x_2, x_4, x_5, y_1, y_2, y_4, y_5),\\
F_4 &= \mathrm{conv}(x_1, x_3, x_4, x_5, y_1, y_3, y_4, y_5),\\
F_5 &= \mathrm{conv}(x_2, x_3, x_4, x_5, y_2, y_3, y_4, y_5),
\end{aligned}$$

where each $\mathrm{conv}(x_i, x_j, x_k, y_i, y_j, y_k)$ is a triangular prism with triangles $\mathrm{conv}(x_i, x_j, x_k)$ and $\mathrm{conv}(y_i, y_j, y_k)$ (we assume that each $\mathrm{conv}(x_k, y_k)$ is an edge of P^*). Using this formula, one conclude that

$$\begin{aligned}
&\partial\langle F_1, F_2, \ldots, F_5\rangle\\
&= \langle \mathrm{conv}(S) : S \subset \{x_1, \ldots, x_5\},\ |S| = 4\rangle \cup \langle \mathrm{conv}(S) : S \subset \{y_1, \ldots, y_5\},\ |S| = 4\rangle
\end{aligned}$$

is the disjoint union of two copies of the boundary of a 4-simplex.

If $\mathrm{conv}(x_1, \ldots, x_5)$ is not a face of P^*, then, for each $S \subset \{x_1, \ldots, x_5\}$ with $|S| = 4$, there is a unique 4-face $G_S \notin \{F_1, \ldots, F_5\}$ of P^* that contains $\mathrm{conv}(S)$. This implies that, since P^* has only 8 facets other than $F_1, \ldots, F_5$, either $\mathrm{conv}(x_1, \ldots, x_5)$ or $\mathrm{conv}(y_1, \ldots, y_5)$ must be a face of P^* (otherwise P^* has at least 10 facets other than $F_1, \ldots, F_5$). We assume that $\mathrm{conv}(x_1, \ldots, x_5)$ is a face of P^*. Let $G_1, \ldots, G_8$ be the simplex 4-faces of P^* and assume $G_1 = \mathrm{conv}(x_1, \ldots, x_5)$. Then $\Gamma = \langle G_2, \ldots, G_8\rangle$ is a pseudomanifold with

$$\partial\Gamma = \langle \mathrm{conv}(S) : S \subset \{y_1, \ldots, y_5\},\ |S| = 4\rangle.$$

Let n_i be the number of interior vertices of Γ. By the Lower Bound Theorem for pseudomanifolds with boundary [Fo] (see also [Ta, Theorem 1.2]), Γ must have at least $5 + 4n_i - 4$ facets. Since Γ only has 7 facets, we have $n_i \leq 1$. However, this implies that Γ is either the 4-simplex or the cone over the boundary of the 4-simplex, contradicting the fact that Γ has 7 facets.

We finally prove that (C) cannot occur. Since $\partial\langle F_1, \ldots, F_5\rangle$ is a simplicial complex, $F_1 \cap F_2$ must be a triangular prism and $F_i \cap F_j$ is a square pyramid for all $i \in \{1, 2\}$ and $j \in \{3, 4, 5\}$. It is not hard to see that $F_1, \ldots, F_5$ can be written as

$$\begin{aligned}
F_1 &= \mathrm{conv}(x, y, z, x', y', z', v_1),\\
F_2 &= \mathrm{conv}(x, y, z, x', y', z', v_2),\\
F_3 &= \mathrm{conv}(x, y, x', y', v_1, v_2),\\
F_4 &= \mathrm{conv}(x, z, x', z', v_1, v_2),\\
F_5 &= \mathrm{conv}(y, z, y', z', v_1, v_2),
\end{aligned}$$

where $\mathrm{conv}(x, y, z, x', y', z')$ is a triangular pyramid with triangles $\mathrm{conv}(x, y, z)$ and $\mathrm{conv}(x', y', z')$ (we assume that $\mathrm{conv}(x, x'), \mathrm{conv}(y, y')$ and $\mathrm{conv}(z, z')$ are edges). A

routine computation shows

$$\begin{aligned}\partial\langle F_1, F_2, \ldots, F_5\rangle =& \langle \mathrm{conv}(S) : S \subset \{v_1, v_2, x, y, z\},\ |S| = 4\rangle \\ & \cup \langle \mathrm{conv}(S) : S \subset \{v_1, v_2, x', y', z'\},\ |S| = 4\rangle\end{aligned}$$

is the union of two copies of the boundary of a 4-simplex intersections in the edge $\mathrm{conv}(v_1, v_2)$. Then the exactly same argument as in the case (b) works, namely, one case show that either $\mathrm{conv}(v_1, v_2, x, y, z)$ or $\mathrm{conv}(v_1, v_2, x', y', z')$ must be a face of P^* and conclude a contradiction by the Lower Bound Theorem for pseudomanifolds with boundary. □

References

[Ba] D.W. Barnette, The projection of the f-vectors of 4-polytopes onto the (E, S)-plane, *Discrete Math.* **10** (1974), 201–209.

[BR] D.W. Barnette and J.R. Reay, Projection of f-vectors of four-polytopes, *J. Combinatorial Theory, Ser. A* **15** (1973), 200–209.

[BL] M. Bayer and C.W. Lee, Combinatorial aspects of convex polytopes, in Handbook of Convex Geometry, P. Gruber and J. Wills, eds., North-Holland, Amsterdam, 1993, pp. 485–534.

[Br] A. Brøndsted, An introduction to convex polytopes, Graduate Texts in Mathematics, vol. 90, Springer, New York, 1983.

[Fo] A. Fogelsanger, The generic rigidity of minimal cycles, Ph.D. Thesis, Cornell University, 1988.

[FMM] K. Fukuda, H. Miyata and S. Moriyama, Classification of Oriented Matroids, abirable at `http://www-imai.is.s.u-tokyo.ac.jp/~hmiyata/oriented_matroids/`

[Gr] B. Grünbaum, Convex Polytopes, Wiley Interscience, New York, 1967.

[Ka] G. Kalai, Rigidity and the lower bound theorem. I, *Invent. Math.* **88** (1987), 125–151.

[PUY] G. Pineda-Villavicencio, J. Ugon and D. Yost, Lower bound theorems for general polytopes, arXiv:1509.08218.

[SZ] H. Sjöberg and G.M. Ziegler, Face and flag vector projections of polytopes, in preparation.

[St] R.P. Stanley, Combinatorics and Commutative Algebra, Second Edition, Birkhäuser, Boston, Basel, Berlin, 1996.

[Ta] T.-S. Tay, Lower-bound theorems for pseudomanifolds, *Discrete Comput. Geom.* **13** (1995), 203–216.

[Zi] G.M. Ziegler, Lectures on polytopes, Graduate Texts in Mathematics, vol. 152, Springer, New York, 1995.

Takuya Kusunoki, Department of Pure and Applied Mathematics, Graduate School of Information Science and Technology, Osaka University, Suita, Osaka, 565-0871, Japan

E-mail address: `t-kusunoki@ist.osaka-u.ac.jp`

Satoshi Murai, Department of Pure and Applied Mathematics, Graduate School of Information Science and Technology, Osaka University, Suita, Osaka, 565-0871, Japan

E-mail address: `s-murai@ist.osaka-u.ac.jp`